

AD-A038 806

EXTRANUCLEAR LABS INC PITTSBURGH PA
CROSS SECTIONS FOR THERMAL REACTIONS BETWEEN URANIUM ATOMS AND --ETC(U)
DEC 76 W L FITE, T A PATTERSON, M W SIEGEL F19628-74-C-0215


F/6 20/8


UNCLASSIFIED

AFGL-TR-77-0030

NL

1 OF
AD
A038806





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 038806

AFGL-TR-77-0030 ✓

(2) 18

CROSS SECTIONS FOR THERMAL REACTIONS BETWEEN URANIUM ATOMS AND ATMOSPHERIC SPECIES

Wade L. Fite
Thomas A. Patterson
Melvin W. Siegel

Extranuclear Laboratories, Inc.
P.O. Box 11512, 250 Alpha Drive
Pittsburgh, PA 15238

December 31, 1976

Final Report for Period 1 November 1974 - 31 December 1976

Approved for Public Release, Distribution Unlimited

This research was sponsored by the Defense Nuclear Agency

AIR FORCE GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AFB, MASSACHUSETTS 01731

AD No. _____
DDC FILE COPY

DDC
RECEIVED
APR 25 1977
B

Qualified requestors may obtain additional copies from the Defense Documentation Center. All others should apply to the National Technical Information Service.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM																															
1. REPORT NUMBER AFGL-TR-77-0030	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9																															
4. TITLE (and Subtitle) CROSS SECTIONS FOR THERMAL REACTIONS BETWEEN URANIUM ATOMS AND ATMOSPHERIC SPECIES.		5. TYPE OF REPORT & PERIOD COVERED Scientific - Final rept. 1 NOV 74 - 31 DEC 76																															
7. AUTHOR(s) W. L. Fite, T. A. Patterson, M. W. Siegel		8. CONTRACT OR GRANT NUMBER(s) F 19628-74-C-0215																															
9. PERFORMING ORGANIZATION NAME AND ADDRESS Extranuclear Laboratories, Inc. P.O. Box 11512 (250 Alpha Drive) Pittsburgh, PA 15238		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE 62704H CDNA-05-AD Subtask S99QAXHD 010, Work Unit 72																															
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory Hanscom AFB, Massachusetts 01730 Contract Monitor: Dr. Edmond Murad/LKB		12. REPORT DATE 31 December 76																															
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Wade L./Fite, Thomas A./Patterson Melvin W./Siegel		13. NUMBER OF PAGES 88																															
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15. SECURITY CLASS. (of this report) Unclassified																															
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 12 87p.																																	
18. SUPPLEMENTARY NOTES This Research was Sponsored by the Defense Nuclear Agency Under Subtask S99QAXHD010, Work Unit 72 Entitled Associative Ionization Reactions of Communication Importance																																	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <table border="0"> <tr> <td>Associative Ionization</td> <td>Ozone</td> <td>Lanthanum</td> <td>SF₄</td> <td>MoO</td> <td>Ozone Source</td> </tr> <tr> <td>Chemi-Ionization</td> <td>Sulfur</td> <td>Gadolinium</td> <td>CHF₃</td> <td>MoO₂</td> <td>Negative ions</td> </tr> <tr> <td>Uranium</td> <td>Thorium</td> <td>Chlorine</td> <td>H₂O₂</td> <td>WO₂</td> <td></td> </tr> <tr> <td>Oxygen</td> <td>Neodymium</td> <td>Bromine</td> <td>(CN)₂</td> <td>TiO</td> <td></td> </tr> <tr> <td>Atomic Oxygen</td> <td>Samarium</td> <td>SF₆</td> <td>H₂S</td> <td>TiO₂</td> <td></td> </tr> </table>				Associative Ionization	Ozone	Lanthanum	SF ₄	MoO	Ozone Source	Chemi-Ionization	Sulfur	Gadolinium	CHF ₃	MoO ₂	Negative ions	Uranium	Thorium	Chlorine	H ₂ O ₂	WO ₂		Oxygen	Neodymium	Bromine	(CN) ₂	TiO		Atomic Oxygen	Samarium	SF ₆	H ₂ S	TiO ₂	
Associative Ionization	Ozone	Lanthanum	SF ₄	MoO	Ozone Source																												
Chemi-Ionization	Sulfur	Gadolinium	CHF ₃	MoO ₂	Negative ions																												
Uranium	Thorium	Chlorine	H ₂ O ₂	WO ₂																													
Oxygen	Neodymium	Bromine	(CN) ₂	TiO																													
Atomic Oxygen	Samarium	SF ₆	H ₂ S	TiO ₂																													
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Experimental studies of associative ionization reactions involving uranium and other metal atoms in collisions with O ₁ O ₂ and other atmospheric species are described. Absolute reaction cross sections are obtained by comparison with the previously measured cross sections for the U + O and U + O ₂ reactions. Negative ions have been observed as products in some of the reactions however these channels account for less than 10% of the total reaction cross section. No reaction products are observed for associative ionization between atomic																																	

388310

20.


oxygen and selected oxides of Mo, W and Ti. Both thorium and uranium have large total cross sections of order 10^{-15} cm², for associative ionization with ozone. A simple, efficient ozone source for beam experiments is described.

10 to the -15 power sq. cm.,

Unclassified

I. Introduction

Associative ionization is that type of chemi-ionization¹ wherein two atoms, molecules or radicals, A and B, collide at thermal energies and produce a molecular ion in a process such as



In order for the process to proceed it is necessary to provide enough energy to remove the electron from the neutral AB molecule or radical, and this energy comes from (1) chemical binding energy of the AB molecule and (2) internal excitation in either A or B.

Examples of associative ionization reactions which rely on internal excitation have long been known, e.g., the formation of diatomic inert gas ions in discharges and afterglows. The molecular ions formed in this situation can rapidly dissociatively recombine with electrons to produce the initial species in their ground states.

In contrast, if sufficient ~~energy~~ for ionization is provided by the formation of the A-B chemical bond, associative ionization can occur between ground state reactants. Neutralization by dissociative recombination is energetically forbidden for molecular ions formed under these conditions and the lifetime of the ions may be long, even in the presence of a substantial electron density. Since the lifetime against neutralization may be long, the possibility exists that the ion, if formed in an excited state, could radiate even if the transition were in the infrared region where excited state lifetimes are long.

Thus reactants that undergo associative ionization from their ground states offer a source of long-lived electron clouds which can affect radar signal transmission and simultaneously a source of infrared radiation which might be unwanted background in any system involving IR radiation.

We began our investigation of associative ionization reactions, under an earlier research contract², by examining the reactions



An absolute measurement of the cross section for the latter reaction³ provided a benchmark for measuring other associative ionization cross sections by a simple comparison technique. Further study led to the discovery that ground state associative ionization occurs in a number of metal-oxygen systems⁴⁻⁸ and a variety of other metal-gas combinations.^{2,4,9}

Much of the effort under the present contract has been devoted to the study of various uranium-gas reactions. The observation of dissociative associative ionization², in particular the production of UF_2^+ from $U + SF_6$, suggested that negative ions might also be formed in some of these reactions. This prompted the development of detection electronics which would permit the observation of negative ions as well as positive ones.

This report summarizes the program as carried out over the past two years, summarizes the present situation with regard to associative ionization processes, and suggests the directions in which future research might profitably move. Included as appendices are a list of associative ionization reactions studied at Extranuclear Laboratories, Inc. and the University of Pittsburgh, abstracts of papers presented at scientific meetings and a preprint of an article on ozone associative ionization with uranium and thorium.

SESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DOC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION.....		
BY.....		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	AND/OR SPECIAL
A		

II. Uranium Associative Ionization With Gases

Much of the work performed under the previous contract to examine associative ionization reactions² was devoted to reactions between atomic and molecular oxygen and various metals. Reactions were observed between uranium and NO, N₂O and SF₆ producing the positive ions UNO⁺, UO⁺ and UF₂⁺, respectively, although detailed quantitative measurements were not performed. With the observation of dissociative associative ionization in the latter two reactions, the possibility arises that negative ions may also be formed in associative ionization reactions.

We first observed the production of negative ions in the U + Br₂ system. Three reaction channels are observed,





By comparison of signal intensities in these reactions with the UO₂⁺ signal from reaction (2) under comparable gas beam intensities, we roughly estimate cross sections of $5 \times 10^{-19} \text{ cm}^2$ for (4) and $1 \times 10^{-19} \text{ cm}^2$ for the sum of (5) and (6). The branching ratio between (5) and (6) is difficult to estimate, but appears to be of order 10:1. The analogous channels are observed in the reactions of Cl₂ with uranium:



with roughly estimated cross sections of 10^{-17} cm^2 for (7) and 10^{-18} cm^2 for the sum of (8) and (9). The branching ratio appears to be of order unity.

Negative ions are also produced in the $\text{U} + \text{SF}_4$ system, where the assumed reactions are



The cross section for (10) is estimated to be $5 \times 10^{-19} \text{ cm}^2$, while the cross section for (11) is so small as to preclude even a rough estimate. The UF_2^+ product in (11) is shown as uncertain because the F^- signal is sufficiently small that a correspondingly small current of a heavy ion, e.g., UF^+ , might easily elude detection. The $\text{U} + \text{SF}_6$ system has a comparable cross section for producing UF_2^+ , however no negative ions are observed.

Since the uranium-atomic oxygen associative ionization reaction has a relatively large cross section, one might expect the analogous reaction with atomic sulfur, which also has an $ns^2 np^4 ({}^3P)$ electron structure, to occur. To investigate this possibility an independently heated solids vaporization stage was added to the existing gas dissociation furnace. With this two-stage furnace, an atomic sulfur beam can be produced by gently vaporizing molecular sulfur in the relatively cool first chamber and dissociating the vapor in the hotter second chamber.

Uranium and sulfur are found to associatively ionize, producing US^+ , an appropriate reaction product to observe during the United States' bicentennial celebration. The reaction cross section is $1.776 \times 10^{-17.76} \text{ cm}^2$, 50 times smaller than the analogous UO cross section.

A cursory examination of the reaction $U + S_2 \rightarrow US_2^+ + e$ yielded a null result. However, considering beam intensities and the ratio of the O and O_2 cross sections, the US_2^+ product could easily have escaped detection. Insufficient data exist on the dissociation energy and ionization potential of US_2 to determine whether the reaction is even energetically possible.

Several other uranium-gas systems have also been investigated, a number of which, CHF_3 , H_2O_2 , $(CN)_2$ and H_2S , produced no observable associative ionization signals. The results of these studies are included in Appendix I.

III. Associative Ionization of Metal Oxides With Atomic Oxygen


Qualitative observations have indicated that certain monoxides such as WO may undergo associative ionization with atomic oxygen. Efforts to confirm this were hampered by physical constraints in the experimental apparatus. There is not sufficient space without extensive modification of the apparatus to replace the simple metal beam furnace with a two-stage oven which would be desirable for producing monoxide and dioxide beams from more highly oxidized metal powders. Consequently we had little control of the oxidation state of the beam. However in experiments with atomic oxygen and MoO, MoO₂, WO₂, TiO and TiO₂, no associative ionization products were observed. Based on experimental conditions, this places an upper bound of $\sim 10^{-18} \text{ cm}^2$ on the cross sections for associative ionization in these systems, if the reaction occur.

IV. Ozone Associative Ionization Reactions With Uranium and Thorium

In associative ionization reactions involving molecular species, multiple reaction channels are conceivable although for most of the systems which we have investigated, insufficient thermodynamic data are available to determine which channels, if any, are energetically allowed. Ozone is one of the few exceptions; thermochemical experiments¹⁰ indicate that five of the seven possible channels are energetically allowed for ozone associative ionization with both uranium and thorium.



We have investigated both the U-O₃ and Th-O₃ systems. Only two of the possible channels are observed in the uranium reaction, but five distinct channels can be identified with thorium. The observed reactions are summarized in Table I. The cross sections, ranging from 10⁻¹⁹ cm² to 10⁻¹⁶ cm², are tabulated in Appendix I; details of the measurements are described in Appendix VIII.

TABLE I
REACTION CHANNELS OBSERVED IN OZONE ASSOCIATIVE
IONIZATION WITH URANIUM AND THORIUM



V. Metal-Oxygen Associative Ionization Reactions

There are three possible associative ionization reaction channels in a metal-molecular oxygen system:



Dr. H. H. Lo at the University of Pittsburgh has observed the production of both MO_2^+ and MO^+ in reactions involving the metals lanthanum and cerium.¹¹ The production of MO^+ implies that the negative ion channels, (14), is at least energetically allowed, although the MO^+ may be due entirely to reaction (13).

We have investigated the La- O_2 system and observe all three reaction channels. The branching ratio between (13) and (14) is estimated to be of order 10:1, yielding a cross section of $\sim 10^{-18} \text{ cm}^2$ for (14) with lanthanum. The cross sections for reactions (12) and (13) with lanthanum are both of order 10^{-17} cm^2 . This is consistent with our general observation that even when negative ions are produced in a reaction, the free electron channels dominate.

We have subsequently learned¹² that Dr. R. B. Cohen and his co-workers at the Illinois Institute of Technology have also observed O^- in the $La + O_2$ reaction and in the $Ce + O_2$ reaction as well.

Lo and Cohen have also each measured cross sections for a number of metal-atomic oxygen associative ionization reactions. As a consistency check on the different experimental systems and techniques, we have examined the neodymium-oxygen and samarium-oxygen reactions which were studied by both groups. Using the previously measured $Gd + O$ reaction

cross section for calibration, we performed direct measurements of the Nd/Gd and Sm/Gd cross section ratios. Setting the ratios of the metal electron impact ionization cross sections equal to unity yields atomic oxygen associative ionization cross sections of $2.7 \times 10^{-15} \text{ cm}^2$ for Nd and $1.7 \times 10^{-16} \text{ cm}^2$ for Sm. A comparison of our results with those of Cohen *et al.* is shown in Table II. While there is some scatter, the two experiments are in reasonable agreement. Lo has also observed these reactions, however his experiments were not intended to produce absolute cross section measurements.

A simultaneous comparison technique is employed in Cohen's experiments. Two different metals are placed in the beam source and the reaction signals are all observed at a constant source temperature. This approach restricts comparisons to metal pairs having comparable vapor pressures in some temperature range. This requires that the Nd-Gd comparison employ a three element chain, Nd/Tb, Tb/Pr, Pr/Gd while two additional elements, Sm/Ho, Ho/Nd, are required for samarium.

TABLE II

COMPARISON OF MEASURED CROSS SECTIONS OF THE REACTION
 $M + O \rightarrow MO^+ + e$ FOR THE METALS NEODYMIUM AND SAMARIUM

	<u>Nd</u>	<u>Sm</u>
Cohen ¹²	$1.3 \times 10^{-15} \text{ cm}^2$	$4.3 \times 10^{-17} \text{ cm}^2$
This Work	$2.7 \times 10^{-15} \text{ cm}^2$	$1.7 \times 10^{-16} \text{ cm}^2$

We have used a sequential comparison method for these experiments. The furnace containing the two metals to be compared is heated slowly and reaction signals are observed as the furnace temperature reaches the appropriate temperature to produce a beam of each metal. Due to the open design of the furnace (described in Appendix III), the lower melting point metal supply is generally exhausted before a beam of the second metal is detected. Thus the signals for the two reactions are measured at different, unknown, furnace temperatures. The geometry of the source also allows a variation in the overlap between the oxygen beam and the two metal beams.

Both techniques have their weak points. We consider our results to be in essential agreement with those of Cohen, and believe that the differences could be reduced by installing a furnace which allows greater control of the metal beam source. For other systems studied by both Lo and Cohen, the agreement is generally satisfactory, less than a factor of ten in most cases.

VI. Technical Developments

In addition to the specific reactions which have been studied, we have developed a number of generally useful experimental techniques. Many of these have been in the area of neutral beam sources. The two-stage furnace which was used to produce an atomic sulfur beam has already been described in Section II. Although the idea of using a two-stage furnace is not new, our experiences have demonstrated that a simple tube-type gas dissociation furnace can be readily adapted to a two-stage design for use with solid samples.

A great deal of effort has been devoted to improving the tungsten foil tube furnace which was used in the early phases of this research project as a uranium beam source. The present design consists of three tungsten rods mounted to form a vee-shaped trough. A chip of uranium is held in the trough by glue which burns away as the furnace is heated, disappearing completely before the uranium begins to vaporize. This furnace has a usable life of five to ten hours with uranium, nearly an order of magnitude longer than the earlier tube furnace.

There are some disadvantages associated with this design. Due to the large physical area over which the beam may be formed, neither the beam geometry nor the beam temperature is well-defined. The source is not well suited for two element beams because the beams of the two metals are produced from different areas of the furnace and are not co-linear.

The ozone beam source design has proved most successful. This all-glass rf discharge source is described in detail in Appendix VIII. The purity of the ozone beam which is produced has not yet been accurately determined due to a lack of data concerning the electron impact ionization cross section and fragmentation pattern of ozone. The source is presently being used, in connection with another project, to measure these parameters.

We have also demonstrated that dc negative ion currents can be routinely measured using an electron multiplier and an Extranuclear Laboratories Negative/Positive Preamplifier. The isolation provided by the preamp allows changing from positive to negative ion detection by simply changing the bias voltages on the ends of the electron multiplier and reversing the polarity of the focusing lens voltages.

VII. Conclusions

The principal conclusion of the present research program is that associative ionization between ground state reactants is not an uncommon process and that it can be a significant source of long-lived electron clouds in the upper atmosphere through the introduction of any of a number of heavy metals. While negative ions may also be produced in some of the reactions, the negative ion channels account for no more than 5% of the total reaction cross section in any of the systems examined thus far.

Knowing the absolute cross sections for the $U + O$ and $U + O_2$ reactions has made it straightforward to obtain cross sections for other associative ionization reactions. These cross section measurements all contain at least one factor which is a ratio of usually poorly known electron impact ionization cross sections, pointing out a need for improved electron impact cross section data. Presently stated results explicitly show this factor so that improved values for these associative ionization cross sections can be generated as soon as improved EI cross sections become available.

It is not yet possible to predict the combinations of reactants which will associatively ionize. Additional data on the ionization potentials and dissociation energies of heavy metal radicals appear to be a necessary, although probably not sufficient, requirement for developing

a predictive capability. Not all of the reaction channels known to be energetically allowed have been observed, however for every combination of reactants known to have one or more allowed channels, at least one channel has been observed.

There remains much to be learned about associative ionization. The possibilities of internal excitation of the product positive ions and the energy distributions of the electrons have not yet been fully examined. There is no doubt as to the basic scientific interest of this entire field and it is quite possible that the processes may also have significant bearing on military communications problems, absolute neutral atom detection, and perhaps isotope separation and other processes in which chemi-ionization between ground state reactants represents an energetically economical means of ionization.

References

1. For a general review of chemi-ionization see A. Fontijn, Progress in Reaction Kinetics, 6, 75 (1972).
2. W. L. Fite, H. H. Lo, P. Irving, M. W. Siegel and R. T. Brackmann, Report No. AFCRL-TR-74-0568, November, 1974.
3. W. L. Fite, H. H. Lo and P. Irving, J. Chem. Phys. 60, 1236 (1974).
4. H. H. Lo and W. L. Fite, Chem. Phys. Lett., 29, 39 (1974).
5. P. D. Zavitsanos, J. Chem. Phys. 59, 2162 (1973).
6. R. B. Cohen, C. E. Young and S. Wexler, Chem. Phys. Lett. 19, 99 (1972).
7. C. E. Young, P. M. Dehmer, R. B. Cohen, L. G. Pobo and S. Wexler, J. Chem. Phys. 64, 306 (1976).
8. C. E. Young, R. B. Cohen, P. M. Dehmer, L. G. Pobo and S. Wexler, J. Chem. Phys. 65, 2562 (1976).
9. R. B. Cohen, P. Majeres and J. K. Raloff, Chem. Phys. Lett 31, 176 (1975).
10. D. L. Hildenbrand, Report No. AFGL-TR-76-0061, March, 1976.
11. H. H. Lo, private communication, 1976.
12. R. B. Cohen, private communication, 1976.

APPENDIX I

ASSOCIATIVE IONIZATION REACTIONS STUDIED AT EXTRANUCLEAR LABORATORIES, INC.
AND THE UNIVERSITY OF PITTSBURGH UP TO DECEMBER 31, 1976

Reaction	Observers	Method	Cross Section (cm ²) See Note 1
1. $U + O \rightarrow UO^+ + e$	1,2,3	A	1.6×10^{-15}
2. $Th + O \rightarrow ThO^+ + e$	1,3	A	1.0×10^{-15}
3. $Zr + O \rightarrow ZrO^+ + e$	1,3	A	1.3×10^{-15}
4. $Gd + O \rightarrow GdO^+ + e$	1,3	A	8.4×10^{-16}
5. $Ti + O \rightarrow TiO^+ + e$	1,3	A	4.5×10^{-15}
6. $La + O \rightarrow LaO^+ + e$	3	A	5.1×10^{-15}
7. $Sm + O \rightarrow SmO^+ + e$	3,6	A	1.7×10^{-16}
8. $Nd + O \rightarrow NdO^+ + e$	3,6	A	2.7×10^{-15}
9. $Y + O \rightarrow YO^+ + e$	3	A	3×10^{-16}
10. $Ce + O \rightarrow CeO^+ + e$	3	A	1×10^{-16}
11. $Pr + O \rightarrow PrO^+ + e$	3	A	2×10^{-15}
12. $Dy + O \rightarrow DyO^+ + e$	3	A	1×10^{-17}
13. $Er + O \rightarrow ErO^+ + e$	3	A	1×10^{-18}
14. $U + O_2 \rightarrow UO_2^+ + e$	1,2,3	A	1.7×10^{-17}
15. $Th + O_2 \rightarrow ThO_2^+ + e$	1,3	A	1.5×10^{-17}
16. $La + O_2 \rightarrow LaO_2^+ + e$	3	A,B	1.7×10^{-17}
17. $\rightarrow LaO^+ + O + e$	3	A,B	1.2×10^{-17}
18. $\rightarrow LaO^+ + O^-$	6	A	10^{-18}
19. $Ce + O_2 \rightarrow CeO_2^+ + e$	3	A	3×10^{-19}
20. $\rightarrow CeO^+ + ?$	3	A	1×10^{-20}

	<u>Reaction</u>	<u>Observers</u>	<u>Method</u>	<u>Cross Section (cm²)</u> <u>See Note 1</u>
21.	$\text{Pr} + \text{O}_2 \rightarrow \text{PrO}_2^+ + \text{e}$	3	A	4×10^{-19}
22.	$\text{U} + \text{O}_3 \rightarrow \text{UO}_2^+ + \text{O} + \text{e}$	5,6	A	2×10^{-16}
23.	$\rightarrow \text{UO}^+ + \text{O}_2 + \text{e}$	5,6	A	2×10^{-16}
24.	$\text{Th} + \text{O}_3 \rightarrow \text{ThO}_3^+ + \text{e}$	5,6	A	2×10^{-19}
25.	$\rightarrow \text{ThO}_2^+ + \text{O} + \text{e}$	5,6	A	1×10^{-17}
26.	$\rightarrow \text{ThO}_2^+ + \text{O}^-$	5,6	A	2×10^{-18}
27.	$\rightarrow \text{ThO}^+ + \text{O}_2 + \text{e}$	5,6	A	4×10^{-16}
28.	$\rightarrow \text{ThO}^+ + \text{O}_2^-$	5,6	A	1×10^{-18}
29.	$\text{U} + \text{Cl}_2 \rightarrow \text{UCl}_2^+ + \text{e}$	5,3	A,B	1×10^{-17}
30.	$\rightarrow \text{UCl}^+ + \text{Cl} + \text{e}$	5,3	A,B	5×10^{-19}
31.	$\rightarrow \text{UCl}^+ + \text{Cl}^- + \text{e}$	5	A	5×10^{-19}
32.	$\text{La} + \text{Cl}_2 \rightarrow \text{LaCl}_2^+ + \text{e}$	3	B	1.1×10^{-16}
33.	$\rightarrow \text{LaCl}^+ + ?$		B	1.2×10^{-16}
34.	$\text{Y} + \text{Cl}_2 \rightarrow \text{YCl}_2^+ + \text{e}$	3	B	9×10^{-18}
35.	$\rightarrow \text{YCl}^+ + ?$	3	B	4×10^{-19}
36.	$\text{U} + \text{Br}_2 \rightarrow \text{UBr}_2^+ + \text{e}$	5	A	5×10^{-19}
37.	$\rightarrow \text{UBr}^+ + \text{Br} + \text{e}$	5	A	1×10^{-19}
38.	$\rightarrow \text{UBr}^+ + \text{Br}^-$	5	A	1×10^{-20}
39.	$\text{U} + \text{NO}_2 \rightarrow \text{UO}_2^+ + \text{N} + \text{e}$	3	B	See Note 2
40.	$\text{U} + \text{N}_2\text{O} \rightarrow \text{UO}^+ + \text{N}_2 + \text{e}$	4,5,3,6	A,B	3×10^{-18} See Note 2
41.	$\text{La} + \text{N}_2\text{O} \rightarrow \text{LaO}^+ + \text{N}_2 + \text{e}$	3	B	1.5×10^{-15}

<u>Reaction</u>		<u>Observers</u>	<u>Method</u>	<u>Cross Section (cm²)</u> <u>See Note 1</u>
42.	$U + NO \rightarrow UNO^+ + e$	4,3	A,B	4.4×10^{-19}
43.	$U + SF_6 \rightarrow UF_2^+ + SF_4 + e$	4,5	A,B	Small
44.	$U + SF_4 \rightarrow UF_2^+ + SF_2 + e$	5	A	5×10^{-19}
45.	$\rightarrow UF_2^+ + SF + F^-$	5	A	10^{-20}
46.	$U + S \rightarrow US^+ + e$	5,6	A	3×10^{-18}
47.	$Ba + OH \rightarrow BAOH^+ + e$	3	A	Confirming Cohen <u>et al.</u>

Observers:

1. W. L. Fite
2. P. Irving
3. H. H. Lo
4. B. N. Kim
5. M. W. Siegel
6. T. A. Patterson

Methods:

- A. Crossed Beams
- B. Beam through gas or magnetic bottle

Note 1: All of the cross sections must be multiplied by the ratio of the electron impact ionization cross section of the metal to that of uranium. The cross sections for reactions 22-47 must also be multiplied by the ratio of the ionization cross section of the gas to that of O₂.

Note 2: U + N₂O in a crossed beam experiment gave a value of 3×10^{-18} cm². However, the value which came from a magnetic bottle experiment was over 10^{-17} cm². We are now trying to understand this discrepancy in terms of systematic errors, probably in the bottle experiment, since a repeat of the U + N₂O crossed beam experiment by a different observer under different conditions also gave 3×10^{-18} cm². Only the bottle experiment has been used to date for the U + NO₂ reaction so we are reluctant to cite a value at this time.

APPENDIX II

PRESENTED AT THE IX INTERNATIONAL CONFERENCE ON THE
PHYSICS OF ELECTRONIC AND ATOMIC COLLISIONS, SEATTLE, JULY 1975

STUDIES OF POSITIVE AND NEGATIVE ION ASSOCIATIVE IONIZATION IN CROSSED BEAMS OF URANIUM AND N_2O , SF_6 , Br_2 , H_2O_2 , CHF_3 AND O_2^+

M. W. SIEGEL AND W. L. FITE

Extranuclear Laboratories, Inc.
P.O. Box 11512, Pittsburgh, PA 15238, U.S.A.


We have recently surveyed six gas-uranium systems seeking associative ionization processes leading to positive and negative ions. In two cases we have been able to determine estimates for associative ionization cross sections relative to the $1.68 \pm 0.27 \times 10^{-17} \text{ cm}^2$ cross section for UO_2^+ from U and O_2 , previously investigated by Fite *et al.*¹ The apparatus and technique used herein are essentially identical to those described in Ref. 1. The only significant change is the addition of a two port gas handling system to allow rapid interchange of gases for normalization purposes.

Quantitative measurements have been made on the UO^+ from U and N_2O system, where the reaction is



The cross section for this process is estimated at 14% of the cross section for UO_2^+ from U and O_2 . This measurement does not yet include several corrections (e.g., O_2 and N_2O beam intensities by electron bombardment, mass discrimination effects, etc.) which will have to be carefully determined before a precise result can be stated. As would be expected, no negative ions were detected.

In the $U + SF_6$ system the UF_2^+ ion is observed, but no negative ion is found. The process involved is thus presumed to be



Its cross section appears to be small. The UF_2^+ is detected at a convenient signal level when the reaction chamber partial pressure due to the SF_6 beam is of the order of 5×10^{-6} torr. When an O_2 beam producing a pressure increase of 1×10^{-6} torr is mixed with the SF_6 beam, the UO_2^+ signal dominates the UF_2^+ signal by a factor of approximately 7, apparently indicating that the cross section for producing UF_2^+ from U and SF_6 appears to be about 3% of the cross section for producing UO_2^+ from U and O_2 .

In the $U + Br_2$ system the UBr_2^+ ion was observed but the UBr^+ was not. Again, negative ions were sought and none observed.

Positive ions (only) were sought in the $U + H_2O_2$ system, but none were observed. Obtaining this null result required a variation on the usual procedure, necessitated by the inseparable admixture of O_2 with the H_2O_2 beam; thus an associative ionization signal is always observed, and it was at first believed to be due to UO_2^+ (or perhaps UO_2H^+) from the $U + H_2O_2$ reaction. To distinguish signal from H_2O_2 from signal O_2 we used a mass stepping device to switch approximately once per second between UO_2^+ and O_2^+ ; the electron beam was switched on only during the O_2^+ dwell periods. The signals were demultiplexed to sample-and-hold amplifiers, and an x-y plot was made of the UO_2^+ vs. O_2^+ signals as the source pressure of O_2 and $(H_2O_2 + O_2)$ were separately varied. These plots had identical slopes, indicating that all of the UO_2^+ signal was due to the O_2 reaction.

Positive ions (only) were sought in the $U + CHF_3$ (Freon 23) system, but a null result was obtained.

In addition to the above newly investigated systems, negative ions were sought in the $U + O_2$ system; again, as expected a null result was obtained.

The null results in our search for negative ions from N_2O , O_2 , Br_2 , and $SF_6 + U$ are disappointing, especially in the latter two cases. We will have to continue to doubt the proper functioning of the apparatus until such time as some associative ionization negative ion is observed. However the experimental procedure is entirely symmetric for negative and positive ions, and we believe that if any of these systems do lead to negative ions, the cross sections are more than an order of magnitude below the smallest positive ion cross section observed to date.

We anticipate that by conference time we will be able to discuss one or two of these new results in quantitative detail; our search for negative ions will also continue, and we hope to be able to report at least one non-null result.

¹W. L. Fite, H. H. Lo, and P. Irving, J. Chem. Phys. 60, 1236, (1974)

* This research was supported by the Defense Nuclear Agency and monitored by the Air Force Cambridge Research Laboratories under Contract Number F 19628-74-C-0215

APPENDIX III

PRESENTED AT THE AMERICAN SOCIETY FOR MASS SPECTROMETRY
CONFERENCE, HOUSTON, MAY 1975

ASSOCIATIVE IONIZATION AND SECONDARY REACTIONS IN $U + O_2$ AND $U + N_2O$ SYSTEMS

W. L. FITE AND H. H. LO, DEPARTMENT OF PHYSICS,
UNIVERSITY OF PITTSBURGH, AND M. W. SIEGEL,
EXTRANUCLEAR LABORATORIES, INC. PITTSBURGH, PA

Crossed beam experiments have yielded the cross section for $U + N_2O \rightarrow UO^+ + N_2 + e$, from the previously measured cross section for $U + O_2 \rightarrow UO_2^+ + e$. In a second experiment a beam of U atoms crossed a magnetic bottle field where gas was added. Ions formed and trapped in the bottle field were examined. With N_2O , UO_2^+ was also present, arising from the ion-molecule reaction $UO^+ + N_2O \rightarrow UO_2^+ + N_2$. With O_2 , UO_3^+ was also present but arising, we believe from energy considerations, from the associative ionization process $UO^* + O_2 \rightarrow UO_3^+ + e$, with the UO coming from $U + O_2 \rightarrow UO^* + O$. The appearance of UO_3^+ suggests possible errors in the known energetics of the uranium-oxygen system. The experiments are described and the results discussed. Possible uses of magnetically confined plasmas of such ions are considered.

Supported in part by the NSF and in part by the Defense Nuclear Agency through AFCRL

APPENDIX IV

PRESENTED AT THE V INTERNATIONAL CONFERENCE ON
ATOMIC PHYSICS, BERKELEY, JULY 1976

STUDY OF THE ASSOCIATIVE IONIZATION REACTION BETWEEN URANIUM AND SULFUR*

T. A. PATTERSON AND M. W. SIEGEL

Extranuclear Laboratories, Inc., Pittsburgh, PA 15238

We are continuing our survey of associative ionization reactions involving uranium atoms by examining the reaction




Since the analogous reaction with atomic oxygen, which like atomic sulfur has an $ns^2 np^4 \{^3p_2\}$ structure, proceeds with a relatively large cross section,¹ $1.62 \pm 0.41 \times 10^{-15} \text{ cm}^2$, it is reasonable to expect that this reaction would occur.

The experiment is performed in a crossed beams configuration using an apparatus and technique which are essentially identical to those described in Ref. 1. A two-stage oven has been added to the system to produce atomic beams from molecular solids. The solid charge is vaporized in the first stage and then dissociated in the hotter second stage. Generally, a single stage oven hot enough to dissociate a molecule will vaporize the charge too rapidly to produce a usable beam.

The cross section for reaction (1) is measured to be of order $3 \times 10^{-18} \text{ cm}^2$, based on a comparison with the $1.68 \pm 0.27 \times 10^{-17} \text{ cm}^2$ cross section for UO_2^+ from U and O_2 , previously investigated by Fite

et al.¹ This value does not include corrections for such factors as the relative efficiencies of the O₂ and S beam intensity measurements by electron bombardment, detailed examination of the S₂ dissociation fraction and mass discrimination effects. These factors will have to be determined to yield a more accurate result.

A cursory examination of the reaction



yielded a null result. However, considering beam intensities and the ratio of the O and O₂ cross sections, the US₂⁺ product could easily have escaped detection. Insufficient data exists on the dissociation energy and ionization potential of US₂ to determine whether the reaction is even energetically possible.

¹W. L. Fite, H. H. Lo and P. Irving, J. Chem. Phys. 60 1236 (1974)

*This research was supported by the Defense Nuclear Agency and monitored by the Air Force Geophysics Laboratory under Contract Number F 19628-74-C-0215

APPENDIX V


PRESENTED AT THE V INTERNATIONAL CONFERENCE ON
ATOMIC PHYSICS, BERKELEY, JULY 1976

STUDIES OF POSITIVE AND NEGATIVE ION ASSOCIATIVE
IONIZATION IN CROSSED BEAMS OF URANIUM AND
Br₂, Cl₂, NO₂, SF₄, (CN)₂ AND H₂S*


M. W. SIEGEL AND W. L. FITE

Extranuclear Laboratories, Inc., Pittsburgh, PA 15238


In a previous report¹ on our crossed beam studies of associative ionization between uranium atoms and various gases we reported observing the reaction



with a null result for the channels





and




Further investigation under more favorably experimental conditions now indicates that reactions (1b) and (1c) do indeed occur. This appears to be the first reported associative ionization reaction leading to a negative ion product. By comparison of observed signal intensities in these cases with the UO_2^+ signal from the reaction $\text{U} + \text{O}_2 \rightarrow \text{UO}_2^+ + e$ under comparable gas beam intensity conditions, we roughly estimate cross sections of $5 \times 10^{-19} \text{ cm}^2$ for (1a), and $1 \times 10^{-19} \text{ cm}^2$ for the sum of (1b) and (1c). The branching ratio between (1b) and (1c) is difficult to estimate, but appears to be the order of 10:1.

We have similarly investigated the reactions of Cl_2 with uranium and observed




and




with roughly estimated cross sections of 10^{-17} cm^2 for (2a), and 10^{-18} cm^2 for the sum of (2b) and (2c). The branching ratio between (2b) and (2c) is, as in the analogous bromine case, difficult to estimate, but appears to be of order unity.

The reaction




previously reported by Fite, Lo and Vasu² in a magnetic bottle configuration, has now been observed in the crossed beam configuration. Its cross section is estimated at $3 \times 10^{-18} \text{ cm}^2$. No negative ion channel is possible.

The reactions



and



have been observed, in interesting contrast to the previously reported¹ reaction $\text{U} + \text{SF}_6 \rightarrow \text{UF}_2^+ + \text{SF}_4 + \text{e}$ where no negative ion channel is observed. The cross section for (4a) is estimated at $5 \times 10^{-19} \text{ cm}^2$,

while the cross section for (4b) is so small as to preclude even a rough estimate at this time. The positive ion product in (4b) is shown as uncertain because the F^- signal is sufficiently small that a correspondingly small current of a heavy positive ion, e.g., UF^+ , might easily elude detection.

The systems $U + (CN)_2$ and $U + H_2S$ were examined and not found to yield ionized products.

References

1. M. W. Siegel and W. L. Fite, Abstracts of Papers of the IXth International Conference on the Physics of Electronic and Atomic Collisions (University of Washington Press, 1975), J. S. Risley and R. Geballe (Eds)., p. 1088.
2. W. L. Fite, H. H. Lo and P. Vasu, Ibid., p. 1086.

*This research was supported by the Defense Nuclear Agency and monitored by the Air Force Geophysics Laboratory under Contract Number F 19628-74-C-0215.

APPENDIX VI

PRESENTED AT THE APS DIVISION OF ELECTRON AND
ATOMIC PHYSICS MEETING, LINCOLN, DECEMBER 1976

OZONE ASSOCIATIVE IONIZATION WITH THORIUM AND URANIUM*

T. A. PATTERSON AND M. W. SIEGEL

Extranuclear Laboratories, Inc., Pittsburgh, PA 15238

Associative ionization reactions between ozone and both uranium and thorium have been studied in a crossed beams experiment. The positive ion products MO_3^+ , MO_2^+ , MO_2^+ and MO^+ , the neutral products O and O_2 , and the negatively charged products e , O^- and O_2^- group into seven associative ionization channels. In the Th-O_3 system all five ion products are observed, indicating that at least three channels are open. Relative intensities of positive and negative ion signals indicate that at least five channels are actually open. The channels leading to $(\text{ThO}^+ + \text{O} + \text{O} + \text{e})$ and $(\text{ThO}^+ + \text{O} + \text{O}^-)$ can probably be excluded on energetic grounds, so cross sections for the five distinct channels can be obtained from the data. In the U-O_3 system, UO^+ and UO_2^+ but not UO_3^+ are observed, in agreement with known energetics. Neither O_2^- nor O^- are observed, although energetically allowed. Measured cross sections span the range 10^{-19} to 10^{-16} cm^2 .

* Supported by the Defense Nuclear Agency and monitored by the Air Force Geophysics Laboratory under Contract Number F 19628-74-C-0215

APPENDIX VII

PRESENTED AT THE APS DIVISION OF ELECTRON AND
ATOMIC PHYSICS MEETING, LINCOLN, DECEMBER 1976

LANTHANUM-OXYGEN ASSOCIATIVE IONIZATION*

T. A. PATTERSON


Extranuclear Laboratories, Inc., Pittsburgh, PA 15238

AND

H. H. LO

University of Pittsburgh, PA 15260

Multiple reaction channels have been observed in heavy metal-molecular halide associative ionization reactions.¹ We report the first observation of multiple channels in a metal-molecular oxygen system. Associative ionization between lanthanum and oxygen has been studied in a crossed beams experiment. In addition to the reaction



which has been observed in many similar metal-oxygen systems,² LaO^+ and O^- are also produced. The cross sections for the first two reactions are of order 10^{-17} cm^2 , while that of the third is significantly smaller.

*Supported by the Defense Nuclear Agency and monitored by the Air Force Geophysics Laboratory under Contract Number F 19628-74-C-0215 and NSF

¹M. W. Siegel and W. L. Fite, Abstracts of the V ICAP (Berkeley, CA, 1976), R. Marrus, M. H. Prior and H. A. Shugart (Eds.), p. 135.

²H. H. Lo and W. L. Fite, Chem. Phys. Lett. 29, 39 (1974) and references therein.

APPENDIX VIII

ASSOCIATIVE IONIZATION OF URANIUM AND THORIUM IN COLLISIONS WITH OZONE*

T. A. PATTERSON, M. W. SIEGEL AND W. L. FITE

Extranuclear Laboratories, Inc., Pittsburgh, PA 15238

ABSTRACT

Associative ionization cross sections for U and Th with O_3 have been measured using a crossed beams technique. Thermochemical data indicate five energetically allowed reaction channels for each metal. Only two of these channels are observed in the U + O_3 system. All five channels, including two leading to the production of negative ions, are observed in the Th- O_3 system. The total reaction cross section for both metals is about $4 \times 10^{-16} \text{ cm}^2$, with individual channel cross sections spanning the range $2 \times 10^{-19} \text{ cm}^2$ to $4 \times 10^{-16} \text{ cm}^2$.

*This research was sponsored by the Defense Nuclear Agency and monitored by the Air Force Geophysics Laboratory under Contract Number F 19628-74-C-0215


ASSOCIATIVE IONIZATION OF URANIUM AND THORIUM
IN COLLISIONS WITH OZONE

T. A. PATTERSON, M. W. SIEGEL AND W. L. FITE

Extranuclear Laboratories, Inc., Pittsburgh, PA 15238



I. Introduction

Associative ionization reactions between lanthanide and actinide heavy metals and a variety of atomic and simple molecular species have been investigated recently.¹⁻⁸ Metal-atom associative reactions have the simple form



Metal-molecule systems are more complex because multiple dissociative-associative reactions channels, including some leading to negative ions, are possible in addition to the simple associative reactions. For many of the heavy metal systems, insufficient molecular binding energy data are available to predict which, if any, of the associative ionization channels are energetically allowed. Turning this deficiency around, the existence or absence of open associative ionization channels may be used to place bounds on certain ionization potentials and dissociation energies.

In this paper we report several associative ionization reactions between ozone and uranium and thorium. Seven associative ionization reaction channels are in principle possible:



Reactions (2)-(6) are known to be exothermic for both uranium⁹ and thorium¹⁰ while (7) and (8) are known to be endothermic by more than 1.5 eV for both metals. Table I summarizes the exothermicity of the reactions. Exothermicity provides only a measure of whether the reaction is permitted, but does not guarantee that the reaction proceeds with sufficient cross section to be observable. Experience has indicated that most but not all allowed reactions are in fact observed. This would be expected on the simple grounds that an enormous number of low lying molecular states exist in these systems, almost assuring one or more appropriate curve crossings in each case.

II. Apparatus

The experiments are performed in the crossed beams apparatus shown schematically in Figure 1. The metal vapor beam and the modulated ozone beam intersect at right angles in the analyzer chamber of a two-chamber, differentially pumped vacuum system. Ions produced by associative

ionization are electrostatically extracted from the interaction volume and focused into the entrance of a quadrupole mass filter. Ions of a selected m/e are detected with a continuous dynode (Channeltron^(R)) multiplier. An electron beam can also be turned on to pass through the interaction volume at right angles to the gas and metal beams. The beam intensities can thus be measured by electron bombardment ionization mass spectrometry.

The metal vapor beam is formed by placing a chip of the metal in a horizontal trough formed by three tungsten rods. The rods are resistively heated to a normal operating temperature of 2000-2200 K. There are three collimating apertures between the beam source and the interaction volume. This very simple technique has been found to be a more satisfactory beam source than the tungsten foil tube furnaces which were used in earlier experiments.¹ The source has an operating life of 8-12 hours with uranium and can easily be replenished many times when used with a less reactive metal.

Ozone is produced in an rf discharge in flowing oxygen at a pressure of 1-10 Torr. The all-glass and teflon ozone source is illustrated in Figure 2. In order to keep O and O₂ discharge products out of the beam and to eliminate electrical noise interference, the ozone is produced and released cyclically. The ozone produced in the discharge is first collected as a solid by cooling the discharge bulb in liquid nitrogen. After collecting approximately .5 cm³ of solid ozone, the discharge and oxygen

flow are turned off. The ozone beam is produced by allowing the solid ozone to evaporate. The ozone effuses from the nozzle of the source into the source chamber of the vacuum system. The nozzle is approximately 1 cm from the beam defining aperture which connects the source chamber with the analyzer chamber.

One of the difficulties in working with a beam of an unstable species such as ozone is to assure that the beam survives the transit from the source to the interaction region. This is particularly important in this case since the ozone dissociation product impurities, O_2 and possibly O , will also react with the metal beam, producing some of the same ions which are expected in the metal-ozone reaction. The only method available for monitoring the ozone beam is to observe the ions produced by electron impact ionization. This introduces a problem in that the ions produced from the impurities by electron impact are also ozone dissociative ionization products. Thus a large O_2^+ signal is not necessarily an indication of an O_2 impurity in the beam. Because the electron impact ionization cross section and fragmentation pattern of ozone is not known, it is not possible to determine the relative concentrations of O , O_2 and O_3 in the beam by observing the ratios of the O^+ , O_2^+ and O_3^+ ion signals. Mass discrimination effects, to be discussed later, would complicate this approach even if the necessary cross section data were available.

Modulating the beam with a rotating toothed wheel permits some information about the purity of the beam to be extracted. By using a frequency selective detection system tuned to the modulation frequency, the

ion signals produced from beam components can be separated from ion signals produced from the background gas. If the phases of the ion signals, relative to some reference signal generated by the modulator, are also monitored, information about the parent neutral(s) may also be obtained, a technique known as phase spectrometry.¹¹

Phase spectrometry is a form of time-of-flight beam analysis. Since neutrals of different masses coming from a source at a given temperature have different velocities, they will have different times of flight in traversing the distance between the modulator and the ionization region. Hence the signal phase of an ion formed by a simple ionization of a neutral will be different from the phase of the same ion formed by dissociative ionization of a more complex neutral. The phase of a particular ion signal depends on the total time of flight, i.e., the neutrals' flight time from the modulator to the ionization region plus the ions' time from the ionization region to the detector. Although the phase is not quite a linear function of the time of flight characteristic of a molecule with the mean velocity, it can be calculated¹² using the path lengths, the source temperature and the ion energy.


The ion signals are treated as two dimensional vectors characterized by an amplitude and a phase. The expected phases of O_3^+ from O_3 , O_2^+ from O_3 and O_2 , and O^+ from O_3 , O_2 and O are calculated and compared with the observed signal phases. In principle, the observed signal vector of

an ion can be resolved into components along the possible source phase axes. This information, together with the relative cross sections for the various ionization reactions, yields a complete description of the neutrals present in the beam.

In practice, the requirement that the ozone beam path length be kept short to maximize the beam intensity in the interaction region leads to the situation that the uncertainty of a phase measurement, typically $\pm 1-2^\circ$, is 10-20% of the phase difference to be detected. The observed signal phases indicate that there are probably small amounts of O_2 in the ozone beam, but without increased phase separation and knowledge of the relative ionization cross sections, it is difficult to make estimates of the impurity concentrations.

III. Procedure

Absolute cross sections are obtained by comparing the product ion current with the product ion current for a similar reaction whose absolute cross section has been carefully measured in another apparatus. The uranium³ and thorium⁴ associative ionization reactions with oxygen,



whose cross sections are $(1.68 \pm 0.27) \times 10^{-17} \text{ cm}^2$ and $1.5 \times 10^{-17} \text{ cm}^2$ respectively, are used as the calibration reactions. The effective associative ionization cross section is proportional to the resultant ion current divided by the product of the ion currents produced by electron impact ionization of the neutral reactant beams.

In practice an oxygen beam is first generated by flowing O_2 through the ozone source. The electron beam is turned on and the modulated O_2^+ and O^+ (from dissociative ionization) signals and the unmodulated M^+ signal are recorded. The electron beam is then turned off and the modulated MO_2^+ associative ionization signal is recorded. The phases of the modulated signals are also noted for later consistency checks.

Ozone is then generated and collected. Using the collected solid ozone at liquid nitrogen temperature as the beam source, the unmodulated M^+ signal and the modulated ozone signals, O_3^+ , O_2^+ and O^+ , produced with the electron beam on, are recorded. The electron beam is then turned off and the observable metal-ozone associative ionization positive ion product signals are recorded. Phases of the modulated signals are again noted. This technique assures reasonably consistent beam overlaps and allows reliable measurement of relative cross sections for the more intense channels.

By removing the liquid nitrogen bath from the ozone collection bulb, the ozone beam intensity can be increased, allowing observation of ion products from other reaction channels having smaller cross sections. As the solid ozone warms, the gas load presented by the source causes the vacuum system pressure to rapidly approach the upper safe operating limit making it necessary to reimmerse the source in liquid nitrogen. This transient nature of the more intense beam requires a modified data acquisition technique.

The quadrupole mass filter is programmed to alternately transmit two different masses, switching masses at one second intervals. The transmitted ion signal is demultiplexed and displayed as two traces on a chart recorder. A previously observed ion product and, at higher signal gain, the mass corresponding to another possible ion product are monitored as the ozone beam intensity is increased. If a new product is observed, the ratio of the reaction channel cross sections is equal to the ratio of the observed ion signals.

This technique is also used to observe the negative ions, O^- and O_2^- which are produced in the thorium-ozone associative ionization reaction. A simultaneous comparison of positive and negative ion yields is not possible with the present system; however the metal atom beam is sufficiently stable and the ozone pressure buildup sufficiently reproducible that reasonably accurate results can be obtained.

IV. Analysis

The observed associative ionization molecular positive ion signal is given by

$$S_3(MO_i^+) = K_3 Q_3(MO_i^+) I_3 n_3 \quad (10)$$

where $Q_3(MO_i^+)$ is the effective cross section for the production of MO_i^+ ($i = 1, 2$ or 3) in the $M-O_3$ reaction. I_3 and n_3 are, respectively the neutral ozone current and the metal number density in the neutral beams. The apparatus constant, K_3 , includes the overall detection efficiency for the product ions and overlap integral of the neutral beams. Similarly, the calibration reaction ion signal can be expressed as

$$S_2(MO_2^+) = K_2 Q_2(MO_2^+) I_2 n_2. \quad (11)$$

The apparatus constants K_3 and K_2 are assumed to be equal. The ion detection efficiency includes the efficiency with which the ions are drawn into the mass filter, the transmission efficiency for the ions and the efficiency of detection at the electron multiplier. Mass discrimination effects in the ion detection can be neglected because the differences in product ion masses are small. Since the source geometry is the same for the oxygen and ozone beams and the metal atom beam is relatively stable, the overlap integrals are approximately equal.

Combining Equations (10) and (11), the ratio of the effective cross section is

$$\frac{Q_3(MO_i^+)}{Q_2(MO_2^+)} = \frac{S_3(MO_i^+)}{S_2(MO_2^+)} \frac{n_2}{n_3} \frac{I_2}{I_3}. \quad (12)$$

The first term is simply the ratio of the observed ion signals. The second term is equal to the ratio of the electron impact ionization metal ion signals recorded during the data runs. This ratio corrects for the small drift in metal beam intensity during the course of the measurements. The third term is the ratio of the O_2 and O_3 neutral beam currents which can be expressed in terms of the number densities and mean molecular speeds as

$$\frac{I_2}{I_3} = \frac{n_2}{n_3} \frac{\bar{v}_2}{\bar{v}_3}. \quad (13)$$

By considering the source temperature and pressure and the length and diameter of room temperature tubing connections the source to the vacuum chamber, it can be shown that the ozone is thermalized from liquid nitrogen temperature to room temperature by the time it enters the beam. Thus the mean molecular speeds are inversely proportional to the square roots of the masses, yielding

$$\frac{I_2}{I_3} = \frac{n_2}{n_3} \left(\frac{M_3}{M_2} \right)^{1/2} = \sqrt{\frac{3}{2}} \frac{n_2}{n_3}. \quad (14)$$

When the ion optics are optimized for maximum sensitivity to the higher mass reaction product ions, significant mass dependent variations in ion transmission may occur at low masses. To avoid transmission factors which are difficult to evaluate, both n_2 and n_3 are expressed in terms of O_2^+ ion signals. Assuming that the ozone beam contains no O_2 , the ratio of the neutral beam currents becomes

$$\frac{I_2}{I_3} = 1.2 \frac{S_2(O_2^+) Q_3^e(O_2^+)}{Q_2^e(O_2^+) S_3(O_2^+)} \quad (15)$$

where $Q_2^e(O_2^+)$ and $Q_3^e(O_2^+)$ are the cross sections for the production of O_2^+ by electron impact on O_2 and O_3 , respectively.

Combining Equations (12), (14) and (15), the expression for the ratio of the effective associative ionization cross sections reduces to

$$\frac{Q_3(MO_i^+)}{Q_2(MO_2^+)} = 1.2 \frac{S_3(MO_i^+) S_2(M^+) S_2(O_2^+) Q_3^e(O_2^+)}{S_2(MO_2^+) S_3(M^+) S_3(O_2^+) Q_2^e(O_2^+)} \quad (16)$$

All but the last term can be evaluated with the experimental data.

Dissociative ionization cross sections for ozone are not presently available.

For future reference when this cross section becomes available, the electron

energy was 50 eV in these experiments. The electron energy for the ratio of the metal ionization cross sections which appears in the thorium-ozone reaction⁴ results was 100 eV.

V. Results

Two uranium-ozone and five thorium-ozone associative ionization reaction channels are observed. The reaction and the measured cross sections are presented in Table II. The quoted cross sections are based on the observations of the first reaction product. Because electron and neutral products are not monitored, the reactions are balanced with the most probable additional products.

The run-to-run scatter in the data indicates an experimental uncertainty of ± 30 -40%. This is small compared with the estimated errors associated with the assumptions made in the interpretation of the data. Assuming equal apparatus constants for the O_2 and O_3 reactions introduces no more than a 5% uncertainty. The most significant assumption is that the ozone beam contains no O_2 or O .

While phase spectroscopy provides some support for this assumption, apparatus constraints limit the sensitivity of this technique. The sign and the magnitude of the effects of 5-10% O and 10-20% O_2 impurity concentrations depend on the reaction product, the reaction cross section and the cross sections for the $M-O$ and $M-O_2$ reactions. These effects can produce changes of a factor of 2-4 in the positive ion cross sections in Table II.

The thorium negative ion cross sections are, at best, order of magnitude estimates. Since the negative ion products are observable only in the transient ozone beam mode, simultaneous comparisons with positive ion signals are not possible. The cross sections are based on comparing the negative ion signals with positive ion signals obtained at another time. In addition to uncertainties associated with transmission factors for positive and negative ions which differ in mass by an order of magnitude, the run-to-run reproducibility of the transient ozone beam is difficult to monitor. A conservative interpretation of the data would be that they do no more than confirm that the reactions occur. Because of this large uncertainty, no attempt has been made to subtract the negative ion contributions to the third and fourth reactions.

VI. Conclusions

There are no apparent correlations between the individual reaction channel cross sections and product ion complexity or reaction exothermicity. The total ozone associative ionization cross sections for both thorium and uranium are of order 10^{-16} cm^2 . These can be compared with atomic oxygen cross sections which are $\sim 10^{-15} \text{ cm}^2$ and molecular oxygen cross sections, $\sim 10^{-17} \text{ cm}^2$. As with previous systems in which negative ion products are observed, the thorium-ozone negative ion channels appear to account for less than 1% of the total cross section. The major negatively charged associative ionization product is a free electron.

1. W. L. Fite and P. Irving, J. Chem. Phys. 56, 4227 (1972)
2. P. D. Zavitsanos, J. Chem. Phys. 59, 2162 (1973)
3. W. L. Fite, H. H. Lo and P. Irving, J. Chem. Phys. 60, 1236 (1974)
4. H. H. Lo and W. L. Fite, Chem. Phys. Lett. 29, 39 (1974)
5. R. B. Cohen, C. E. Young and S. Wexler, Chem. Phys. Lett. 19, 99 (1972)
6. C. E. Young, P. M. Dehmer, R. B. Cohen, L. G. Pobo and S. Wexler, J. Chem. Phys. 64, 306 (1976)
7. C. E. Young, R. B. Cohen, P. M. Dehmer, L. G. Pobo and S. Wexler, J. Chem. Phys. 65, 2562 (1976)
8. R. B. Cohen, private communication, 1976
9. D. L. Hildenbrand, AFGL-TR-76-0061 (1976)
10. D. L. Hildenbrand and E. Murad, J. Chem. Phys. 61, 1232 (1974)
11. W. L. Fite, International Journal of Mass Spectrometry and Ion Physics 16, 109 (1975)
12. H. Harrison, D. G. Hummer and W. L. Fite, J. Chem. Phys. 41, 2567 (1964)

TABLE I
REACTION EXOTHERMICITY (eV)

	<u>U</u>	<u>Th</u>
$M + O_3 \rightarrow MO_3^+$	+ 4.9	≥ 1.8
$\rightarrow MO_2^+ + O + e$	+ 3.2	+ 1.8
$\rightarrow MO_2^+ + O^-$	+ 4.7	+ 3.3
$\rightarrow MO^+ + O_2 + e$	+ 1.1	≤ 1.8
$\rightarrow MO^+ + O_2^-$	+ 2.6	+ 2.3
$\rightarrow MO^+ + O + O + e$	- 4.0	- 3.3
$\rightarrow MO^+ + O + O^-$	- 2.5	- 1.8

TABLE II
REACTION CROSS SECTIONS

<u>Reaction</u>	<u>Cross Section (cm²)</u> See Note 1
U + O ₃ → UO ₂ ⁺ + O + e	2 x 10 ⁻¹⁶
→ UO ⁺ + O ₂ + e	2 x 10 ⁻¹⁶
Th + O ₃ → ThO ₃ ⁺ + e	2 x 10 ⁻¹⁹
→ ThO ₂ ⁺ + O + e	1 x 10 ⁻¹⁷
→ ThO ⁺ + O ₂ + e	4 x 10 ⁻¹⁶
→ O ₂ ⁻ + ThO ⁺	1 x 10 ⁻¹⁸
→ O ⁻ + ThO ₂ ⁺	2 x 10 ⁻¹⁸

Note 1

All cross sections must be multiplied by the ratio of the cross sections for the production of O₂⁺ by electron impact on O₃ and O₂. The thorium cross sections must also be multiplied by the ratio of the thorium and uranium electron impact ionization cross sections.

FIGURE CAPTIONS

Figure 1. Crossed Beams Experimental Arrangement

Figure 2. Ozone Source

FIGURE 1





FIGURE 2



083

DATE 1/02/77

DEPARTMENT OF DEFENSE

DISTRIBUTION LIST

DIRECTOR
DEFENSE ADVANCED RSCH PROJ AGENCY
ARCHITECT BUILDING
1400 WILSON BLVD.
ARLINGTON, VA 22209

01CY ATTN STO KENT KRESA
01CY ATTN LTC W A WHITAKER
01CY ATTN STO CAPT J JUSTICE
01CY ATTN TID FRED A KOETHER

DEFENSE DOCUMENTATION CENTER
CAMERON STATION
ALEXANDRIA, VA 22314

(12 COPIES IF OPEN PUBLICATION, OTHERWISE 2 COPIES)
12CY ATTN TC

DIRECTOR
DEFENSE INTELLIGENCE AGENCY
WASHINGTON, DC 20301
01CY ATTN DT-1 MR KNOLL
01CY ATTN DT-1C

DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON, DC 20305
03CY ATTN STTL TECH LIBRARY
01CY ATTN RAAF P FLEMING
03CY ATTN RAAF CHARLES A BLANK
01CY ATTN STSI ARCHIVES
01CY ATTN STRA
01CY ATTN RAAF MAJ JOHN CLARK
01CY ATTN RAAF HAROLD C FITZ JR

PRECEDING PAGE, BLANK, NOT

083

DATE 23/02/77

DEPARTMENT OF DEFENSE

COMMANDER
FIELD COMMAND
DEFENSE NUCLEAR AGENCY

KIRTLAND AFB, NM 87115

OICY ATTN FCT WILLIAM S ISENGARD

OICY ATTN FCPD

DIRECTOR
INTERSERVICE NUCLEAR WEAPONS SCHOOL

KIRTLAND AFB, NM 87115

OICY ATTN DOCUMENT CONTROL

DIRECTOR
JOINT STRAT TGT PLANNING STAFF JCS
OFFUTT AFB

OMAHA, NE 68113

OICY ATTN JPST MAJ J S GREEN

CHIEF
LIVERMORE DIVISION FLD COMMAND DNA
LAWRENCE LIVERMORE LABORATORY

P.O. BOX 808

LIVERMORE, CA 94550

OICY ATTN FCPRL

083

DATE 1/02/77

DEPARTMENT OF ARMY

COMMANDER/DIRECTOR
ATMOSPHERIC SCIENCES LABORATORY
U S ARMY ELECTRONICS COMMAND
WHITE SANDS MISSILE RANGE, NM 88002

01CY ATTN E BUTTERFIELD DRSEL-BL-SY-R
01CY ATTN R OLSEN
01CY ATTN M DIAMOND
01CY ATTN H BALLARD
01CY ATTN DRSEL-BL-SY-S F E NILES

DIRECTOR
RMD ADVANCED TECH CTR
HUNTSVILLE OFFICE
P.O. BOX 1500

HUNTSVILLE, AL 35807
01CY ATTN ATC-T MELVIN T CAPPS
01CY ATTN ATC-D W DAVIES

CHIEF C-E SERVICES DIVISION
US ARMY COMMUNICATIONS CMD
PENTAGON RM 1R269
WASHINGTON, DC 20310
01CY ATTN CC-OPS-CE

DEP CHIEF OF STAFF FOR RSCH DEV & ACQ
DEPARTMENT OF THE ARMY
WASHINGTON, DC 20310
01CY ATTN ARMMA-00
01CY ATTN SAFEGUARD AND SPACE DIV

083

DATE 23/02/77

DEPARTMENT OF ARMY

COMMANDER

HARRY DIAMOND LABORATORIES
2800 POWDER MILL ROAD
ADELPHI MD 20783

(CNWDI-INNER ENVELOPE: ATTN: DRXDD-RBH)

02CY ATTN DRXDD-NP

01CY ATTN MILDRED H WEINER DRXDD-TT

COMMANDER

TRASANA
WHITE SANDS MISSILE RANGE NM 88002
01CY ATTN R E DEKINDER JR

DIRECTOR

U S ARMY BALLISTIC RESEARCH LABS
ABERDEEN PROVING GROUND, MD 21005

01CY ATTN DRXRR-AM D SNIDER

01CY ATTN J A VANDERHOFF

01CY ATTN DRXRR-CA T J BRIEN

01CY ATTN DRXRR-AM J S RATTEN

01CY ATTN T L CHIDSEY

01CY ATTN DRXRR-AM G E KELLER

01CY ATTN J HEIMERL DRXRD-BSP

01CY ATTN JOHN C MESTER

01CY ATTN MARK D KREGEL

01CY ATTN TECH LIE EDWARD BAICY

COMMANDER

U S ARMY ELECTRONICS COMMAND
FORT MONMOUTH, NJ 07703

01CY ATTN DRSEL-PL-ENV HANS A BOMKE

083

DATE 1/02/77

DEPARTMENT OF ARMY

COMMANDER
II S ARMY FOREIGN SCIENCE & TECH CTR
220 7TH STREET, NE
CHARLOTTESVILLE, VA 22901
OICY ATTN R JONES

COMMANDER
II S ARMY MATERIEL DEV & READINESS CMD
5001 EISENHOWER AVENUE
ALEXANDRIA, VA 22333
OICY ATTN DRCD-0 LAWRENCE FLYNN

COMMANDER
II S ARMY NUCLEAR AGENCY
FORT BLISS, TX 79916
OICY ATTN MONA-WE J BERRERET

CHIEF
IIS ARMY RESEARCH OFFICE
P.O. BOX 12211
TRIANGLE PARK, NC 27709
OICY ATTN CRDARD-P ROBERT MACE

083

DATE 23/02/77

DEPARTMENT OF NAVY

CHIEF OF NAVAL OPERATIONS

NAVY DEPARTMENT

WASHINGTON, DC 20350

OICY ATTN OP 085

CHIEF OF NAVAL RESEARCH

NAVY DEPARTMENT

ARLINGTON, VA 22217

OICY ATTN CODE 464 THOMAS P QUINN

OICY ATTN CODE 464 R GRACEN JOINER

OICY ATTN CODE JTH JOHN G DARDIS

OICY ATTN CODE 427 CDR RONALD J ORERLE

COMMANDER

NAVAL ELECTRONICS LABORATORY CENTER

SAN DIEGO, CA 92152

OICY ATTN WILLIAM F MOLER

OICY ATTN CODE 2200 JURGEN RICHTER

OICY ATTN CODE 2200 HERRERT HUGHES

OICY ATTN CODE 2200 RICHARD PAPPERT

OICY ATTN CODE 2200 ILAN ROTHMILLER

SUPERINTENDENT (CODE 1424)

NAVAL POSTGRADUATE SCHOOL

MONTEREY, CA 93940

(DESIRES NO CNWDI DOCUMENTS)

OICY ATTN CODE 61 MN E A MTLEF

DIRECTOR

NAVAL RESEARCH LABORATORY

WASHINGTON, DC 20375

OICY ATTN CODE 7701 JACK D BROWN

OICY ATTN CODE 7127 CHARLES Y JOHNSON

083

DATE 1/02/77

DEPARTMENT OF NAVY

01CY ATTN CODE 7750 J DAVIS

01CY ATTN CODE 7750 WAHAB ALI
TIMOTHY P COFFEY

01CY ATTN CODE 7750 KLAUS HAIN

01CY ATTN CODE 7750 PAUL JULIENNE

01CY ATTN CODE 7730 EDGAR A MCCLENN

01CY ATTN CODE 2027 TECH LIR

01CY ATTN CODE 7750 DARRELL F STORREL

COMMANDER

NAVAL SURFACE WEAPONS CENTER

WHITE OAK, SILVER SPRING, MD 20910

01CY ATTN CODE WA501 NAVY NUIC PRGMS OFF

01CY ATTN I RUDLIN

01CY ATTN D J LAND

083

DATE 23/02/77

DEPARTMENT OF THE AIR FORCE

AF GEOPHYSICS LABORATORY, AFSC
HANSCOM AFB, MA 01731

01CY ATTN LKB A FAYRE
01CY ATTN DPR ALVA T STAIR
01CY ATTN DPR J RODGERS
01CY ATTN LKD ROCCO S NARCISI
01CY ATTN LKB JOHN PAULSON
01CY ATTN LKB L WEEKS
01CY ATTN DPR JAMES C ULWICK
01CY ATTN LKB EDMOND MURAD
01CY ATTN LKB D GOLOMB
01CY ATTN LKB KENNETH S W CHAMPTON
01CY ATTN LKB T J KENESHEA
01CY ATTN LKB WILLIAM SWIDER JR
01CY ATTN LKD C R PHILBRICK
01CY ATTN DP JOHN S GARING
01CY ATTN LKS F R INNES
01CY ATTN LKS R A VAN TASSEL
01CY ATTN LKD ROBERT E HUFFMAN
01CY ATTN DPR J KENNEALY
01CY ATTN DPR F DELEGRECH
01CY ATTN DPR HAROLD GARDNER
01CY ATTN DPR J H SCHUMMERS
01CY ATTN DPR R ONEILL
01CY ATTN DPR R E MURPHY
01CY ATTN SUOI AFCL RSCH LIB

AF WEAPONS LABORATORY, AFSC
KIRTLAND AFB, NM 87117

01CY ATTN DYT CAPT RICHARD A HARRIS
01CY ATTN 01DYT LT JOHN LILLIS
01CY ATTN DYT CAPT DANIEL A MATUSKA
01CY ATTN DYT CAPT GARY D CABLE
01CY ATTN DYT CAPT DAVID W GOFTZ
01CY ATTN DYT CAPT HOLLARS
01CY ATTN DYT JOSEPH JANNT
01CY ATTN SUL

083

DATE 1/02/77

DEPARTMENT OF THE AIR FORCE

AFTAC

PATRICK AFB, FL 32925

01CY ATTN TF/MAJ WILEY

01CY ATTN TF MAJ E HINES

01CY ATTN TN

HEADQUARTERS

AIR FORCE SYSTEMS COMMAND

ANDREWS AFB

WASHINGTON, DC 20331

01CY ATTN SDR

HEADQUARTERS

ELECTRONIC SYSTEMS DIVISION, (AFSC)

HANSCOM AFB, MA 01731

01CY ATTN ESTI

COMMANDER

FOREIGN TECHNOLOGY DIVISION, AFSC

WRIGHT-PATTERSON AFB, OH 45433

01CY ATTN TD-RTA LIBRARY

01CY ATTN MAJ F FREY

HQ USAF/IN

WASHINGTON, DC 20330

01CY ATTN INS

083

DATE 23/02/77

DEPARTMENT OF THE AIR FORCE

HQ USAF/RD

WASHINGTON, DC 20330

OICY ATTN RDO H F KERR

OICY ATTN RD DCS D RAND

OICY ATTN RDQPN

COMMANDER
ROME AIR DEVELOPMENT CENTER, AFSC

GRIFFISS AFB, NY 13440

OICY ATTN V COYNE

OICY ATTN FMTID DDC LIBRARY

OICY ATTN DCSF J J SIMONS

OICY ATTN TOR

SAMSON/SZ

POST OFFICE BOX 92960

WORLDWAY POSTAL CENTER

LOS ANGELES, CA 90009

(SPACE DEFENSE SYSTEMS)

OICY ATTN SZJ MAJOR LAWRENCE DODAN

COMMANDER IN CHIEF
STRATEGIC AIR COMMAND

OFFUTT AFB, NE 68113

OICY ATTN XPFS MAJ BRIAN G STEPHAN

083

DATE 1/02/77

U S ENERGY RSCH AND DEV ADMIN

ARGONNE NATIONAL LABORATORY
RECORDS CONTROL
9700 SOUTH CASS AVENUE
ARGONNE, IL 60439

01CY ATTN DDC CON FOR GERALD T REEDY
01CY ATTN DDC CON FOR LEN LEIRWITZ
01CY ATTN DDC CON FOR DAVID W GREEN
01CY ATTN DDC CON FOR S GABELNYCK
01CY ATTN DDC CON FOR LTB SVCS RPTS SEC
01CY ATTN DDC CON FOR A C WAHL

DIVISION OF MILITARY APPLICATION
U S ENERGY RSCH & DEV ADMIN
WASHINGTON, DC 20545

01CY ATTN DDC CON FOR MAJ D A HAYCOCK

UNIVERSITY OF CALIFORNIA
LAWRENCE LIVERMORE LABORATORY
P.O. BOX 808
LIVERMORE CA 94550

01CY ATTN D J WUERBLES L-142
01CY ATTN W H DUEWER GEN L-404
01CY ATTN G R HAUGEN L-404
01CY ATTN JULIUS CHANG L-71

LOS ALAMOS SCIENTIFIC LABORATORY
P.O. BOX 1663
LOS ALAMOS, NM 87545

01CY ATTN DDC CON FOR JOHN S MALIK
01CY ATTN DDC CON FOR GROUND CMB-T
01CY ATTN DDC CON FOR T RIENIEWSKI CMB-1
01CY ATTN DDC CON FOR HERMAN HOERLIN
01CY ATTN DDC CON FOR J JUDD
01CY ATTN DDC CON FOR JOHN ZINN

083

DATE 23/02/77

U S ENERGY RSCH AND DEV ADMIN

01CY ATTN D M ROHRER MS 518
01CY ATTN DDC CON FOR E A BRYANT (NO DASTAC)
01CY ATTN DDC CON FOR W M HUGHES MS 559
01CY ATTN DDC CON FOR W D BARFIELD
01CY ATTN DDC CON FOR MARGE JOHNSON
01CY ATTN DDC CON FOR WILLIAM MATER
01CY ATTN DDC CON FOR MARTIN TIERNEY J-10
01CY ATTN DDC CON FOR DAVID W STEINHAUS

SANDIA LABORATORIES
P.O. BOX 5800

ALBUQUERQUE, NM 87115

01CY ATTN DDC CON FOR D A DAHLGREN ORG 1722
01CY ATTN DDC CON FOR FRANK P HINDSON ORG 1722
01CY ATTN DDC CON FOR ORG 3422-1 SANDIA RPT COLL
01CY ATTN DDC CON FOR CLARENCE R MEHL ORG 5230

U S ENERGY RSCH & DEV ADMIN
DIVISION OF HEADQUARTERS SERVICES
LIBRARY BRANCH G-043
WASHINGTON, DC 20545

01CY ATTN DDC CON FOR CLASS TECH LTR

083

DATE 1/02/77

OTHER GOVERNMENT

ALBANY METALLURGY RESEARCH CENTER

U.S. BUREAU OF MINES

P.O. BOX 70

ALBANY, OR 97321

01CY ATTN ELEANOR ARSHIRE

CENTRAL INTELLIGENCE AGENCY

ATTN: RD/SI RM 5G48 HQ BLDG

WASHINGTON, DC 20505

01CY ATTN NED/OSI - 2G48 HQS

DEPARTMENT OF COMMERCE

NATIONAL BUREAU OF STANDARDS

WASHINGTON, DC 20234

(ALL CORRES: ATTN SEC OFFICER FOR)

01CY ATTN K KESSLER

01CY ATTN GEORGE A SINNATT

01CY ATTN J COOPER

01CY ATTN STANLEY ABRAMOWITZ

01CY ATTN M KRAUSS

01CY ATTN LEWIS H GEVANTMAN

DEPARTMENT OF COMMERCE

OFFICE OF TELECOMMUNICATIONS

INSTITUTE FOR TELECOM SCIENCE

BOULDER, CO 80302

01CY ATTN L A BERRY

01CY ATTN WILLIAM F UTLAUT

083

DATE 23/02/77

OTHER GOVERNMENT

DEPARTMENT OF TRANSPORTATION
TRANSPORTATION RSCH SYSTEM CENTER
KENDALL SQUARE
CAMBRIDGE, MA 02142
OICY ATTN F MARMO

DEPARTMENT OF TRANSPORTATION
OFFICE OF THE SECRETARY
TAD-44.1, ROOM 10402-B
400 7TH STREET S.W.
WASHINGTON, D.C. 20590
OICY ATTN RICHARD L STROMBOTNE TST-B
OICY ATTN R UNDERWOOD

NASA
GODDARD SPACE FLIGHT CENTER
GREENBELT, MD 20771
OICY ATTN A TEMPKIN
OICY ATTN S J BAUER
OICY ATTN TECHNICAL LIBRARY
OICY ATTN A C ATKEN

NASA
800 INDEPENDENCE AVENUE S W
WASHINGTON, DC 20546
OICY ATTN R FELLOWS
OICY ATTN D P CAUFFMAN
OICY ATTN LTC D R HALLENBECK CODE SG
OICY ATTN A GESSOW

083

DATE 1/02/77

OTHER GOVERNMENT

NASA

LANGLEY RESEARCH CENTER

LANGLEY STATION

HAMPTON, VA 23365

01CY ATTN CHARLES SCHEXNAYDER MS-168

NASA

AMES RESEARCH CENTER

MOFFETT FIELD, CA 94035

01CY ATTN N-254-3 NEIL H FARLOW

01CY ATTN N-254-4 WALTER L STARR

01CY ATTN M-254-4 R WHITTEN UNCL ONLY

01CY ATTN N-245-4 ILIA G POPPOFF

NATIONAL OCEANIC & ATMOSPHERIC ADMIN

ENVIRONMENTAL RESEARCH LABORATORIES

DEPARTMENT OF COMMERCE

BOULDER, CO 80302

01CY ATTN FRED FEHSENFELD

01CY ATTN FLOON FERGUSON

01CY ATTN J W POPE

01CY ATTN GEORGE C REID ASTRONOMY LAB

NATIONAL SCIENCE FOUNDATION

1800 G STREET, N.W.

WASHINGTON, DC 20556

(NO CLASSIFIED TO THIS ADDRESS)

01CY ATTN F P TODD

01CY ATTN R S ZAPOLSKY

01CY ATTN W D ADAMS

01CY ATTN ROLE SINCLAIR

01CY ATTN R MANKA

01CY ATTN FRED D WHITE

083

DATE 23/02/77

OTHER GOVERNMENT

01CY ATTN M K WILSON

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

AERO-CHEM RESEARCH LABORATORIES, INC

P.O. BOX 12

PRINCETON, NJ 08540

01CY ATTN A FONTIJN

01CY ATTN H PERGAMENT

01CY ATTN HARTWELL F CALCHT

AERODYNE RESEARCH, INC.

REDFORD RESEARCH PARK

CROSBY DRIVE

REDFORD, MA 01730

01CY ATTN F BIEN

01CY ATTN M CAMAC

AERONOMY CORPORATION

217 S NEIL STREET

CHAMPAIGN, IL 61820

01CY ATTN S A BOWHILL

AEROSPACE CORPORATION

P.O. BOX 92957

LOS ANGELES, CA 90009

01CY ATTN N COHEN

01CY ATTN R J MCNEAL

01CY ATTN STONEY W KASH

01CY ATTN R GROVE

01CY ATTN T WIDHOPH

01CY ATTN R D RAWCLIFFE

01CY ATTN JULIAN REINHETER

01CY ATTN IRVING M GARFINKEL

01CY ATTN V JOSEPHSON

01CY ATTN HARRIS MAYER

01CY ATTN T TAYLOR

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

~~AVCO-EVERETT RESEARCH LABORATORY INC~~

~~2385 REVERE BEACH PARKWAY
EVERETT, MA 02149~~

01CY ATTN GEORGE SUTTON

01CY ATTN C W VON ROSENBERG JR

01CY ATTN TECHNICAL LIBRARY

~~RATTELLE MEMORIAL INSTITUTE
505 KING AVENUE~~

~~COLUMBUS, OH 43201~~

01CY ATTN RICHARD K THATCHER

01CY ATTN SYDYAC

01CY ATTN DONALD J HAMMAN

~~THE TRUSTEES OF
BOSTON COLLEGE~~

~~CHESTNUT HILL CAMPUS~~

~~CHESTNUT HILL, MA 02167~~

02CY ATTN CHAIRMAN DEPT OF CHEM

~~BROWN ENGINEERING COMPANY, INC.~~

~~CUMMINGS RESEARCH PARK~~

~~HUNTSVILLE, AL 35807~~

01CY ATTN N PASSINO

01CY ATTN RONALD PATRICK

~~CALIFORNIA AT RIVERSIDE, UNIV OF
RIVERSIDE, CA 92502~~

01CY ATTN JAMES N PITTS JR

01CY ATTN ALAN C LLOYD

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

CALIFORNIA AT SAN DIEGO, UNIV OF
BLDG. 500 MATTHEWS CAMPUS
3175 MIRAMAR ROAD
LA JOLLA, CA 92037
OICY ATTN S C LIN

CALIFORNIA INSTITUTE OF TECHNOLOGY
JET PROPULSION LABORATORY
4800 OAK PARK GROVE
PASADENA, CA 91103
OICY ATTN JOSEPH AJELLO

CALIFORNIA UNIVERSITY OF
BERKELEY, CAMPUIS, ROOM 8
2543 CHANNING WAY
BERKELEY, CA 94720

(CLASS MAIL TO: ATTN SEC OFFICER FOR)

OICY ATTN F MOZER
OICY ATTN HAROLD JOHNSTON
OICY ATTN DEPT OF CHEM W H MILLER

CALIFORNIA, STATE OF
AIR RESOURCES BOARD
9528 TELSTAR AVENUE
FL MONTE, CA 91731
OICY ATTN LEO ZAFONTE

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

CALSPAN CORPORATION

P.O. BOX 235

BUFFALO, NY 14221

01CY ATTN ROBERT A FLUEGG

01CY ATTN M G DUNN

01CY ATTN W WURSTER

01CY ATTN C E TREANOR

01CY ATTN G C VALLEY

COLORADO, UNIVERSITY OF
OFFICE OF CONTRACTS AND GRANTS

380 ADMINISTRATIVE ANNEX

BOULDER, CO 80302

01CY ATTN JEFFREY R PEARCE LASP

01CY ATTN C LINERBERGER JILA

01CY ATTN A PHELPS JILA

01CY ATTN C BEATY JILA

01CY ATTN CHARLES A BARTH LASP

COLUMBIA UNIVERSITY, THE TRUSTEES
IN THE CITY OF NEW YORK

LA MONT DOHERTY GEOLOGICAL

OBSERVATORY - TIDDEY CLIFF

PALISADES, NY 19064

01CY ATTN R PHELAN

COLUMBIA UNIVERSITY, THE TRUSTEES OF
CITY OF NEW YORK

116TH STREET & BROADWAY

NEW YORK, NY 10027

01CY ATTN SFC OFFICER H M FOLEY

01CY ATTN RICHARD N ZARE

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

CONCORD SCIENCES

P.O. BOX 119

CONCORD, MA 01742

OICY ATTN EMMETT A SUTTON

DENVER, UNIVERSITY OF

COLORADO SEMINARY

DENVER RESEARCH INSTITUTE

P.O. BOX 10127

DENVER, CO 80210

(ONLY 1 COPY OF CLASS RPTS)

OICY ATTN SEC OFFICER FOR MR VAN ZYL

OICY ATTN SEC OFFICER FOR DAVID MURCRAE

ECOMETRICS, INC.

718 WILLOW GLEN ROAD

SANTA BARBARA, CA 93105

OICY ATTN NYLF G UTTERBACK

EPSILON LABORATORIES, INC.

4 PRESTON COURT

REDFORD, MA 01730

OICY ATTN JOHN DALCHINES

OICY ATTN HENRY MIRANDA

OICY ATTN CARL ACCARDO

FSL, INC.

495 JAVA DRIVE

SUNNYVALE, CA 94086

OICY ATTN JAMES MARSHALL

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

01CY ATTN W R BELI

GENERAL ELECTRIC COMPANY
SPACE DIVISION
VALLEY FORGE SPACE CENTER
GODDARD BLVD KING OF PRUSSIA
P.O. BOX 8555

PHILADELPHIA PA 19101

01CY ATTN T BAURER

01CY ATTN ROBERT H EDSALL

01CY ATTN P ZAVITSANOS

01CY ATTN M H RORTNER SPACE SCI LAB

01CY ATTN F ALYEA

01CY ATTN J BURNS

01CY ATTN M LINEVSKY

GENERAL ELECTRIC COMPANY
TEMPORARY CENTER FOR ADVANCED STUDIES
816 STATE STREET (P.O. DRAWER 00)
SANTA BARBARA, CA 93102

01CY ATTN DASTAC ART FERRYOK

01CY ATTN TIM STEPHENS

01CY ATTN DON CHANDLER

01CY ATTN WARREN S KNAPP

01CY ATTN R GAMBITL

01CY ATTN DASTAC

GENERAL MOTORS CORPORATION
DELCO ELECTRONICS DIVISION
SANTA BARBARA OPERATIONS
6767 HOLLISTER AVENUE
GOLETA, CA 93017

01CY ATTN ROBIN I PRIMICH

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

GENERAL RESEARCH CORPORATION
P.O. BOX 3587

SANTA BARBARA, CA 93105

OICY ATTN H H LEWIS

OICY ATTN JOHN TSE JR

GENERAL RESEARCH CORPORATION
WASHINGTON OPERATIONS

WESTGATE RESEARCH PARK

7655 OLD SPRINGHOUSE ROAD, SUITE 700

MCLEAN, VA 22101

OICY ATTN THOMAS M ZAKRZEWSKI

GEOPHYSICAL INSTITUTE
UNIVERSITY OF ALASKA

FAIRBANKS, AK 99701

(ALL CLASS ATTN: SECURITY OFFICER)

OICY ATTN HENRY COLE

OICY ATTN R PARTHASARATHY

OICY ATTN NEAL BROWN (UNCL ONLY)

OICY ATTN R J WATKINS

OICY ATTN T N DAVIS (UNCL ONLY)

OICY ATTN J S WAGNER PHYSICS DEPT

OICY ATTN D J HENDERSON

GOVT FISCAL RELATIONS & PATENT OFF

275 ADMIN BUILDING AG-70

UNIVERSITY OF WASHINGTON

SEATTLE WA 98195

OICY ATTN KENNETH C CLARK

OICY ATTN R GERALIE

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

HARVARD UNIVERSITY
DEPARTMENT OF ATMOSPHERIC SCIENCES
PIERCE HALL
CAMBRIDGE, MA 02138
OICY ATTN MICHAEL B MCCLROY

HOWARD UNIVERSITY
DEPARTMENT OF CHEMISTRY
WASHINGTON, DC 20059
OICY ATTN WILLIAM JACKSON

HSS, INC.
2 ALFRED CIRCLE
REDFORD, MA 01730
OICY ATTN H STEWART
OICY ATTN M P SHULER
OICY ATTN DONALD HANSEN

IIT RESEARCH INSTITUTE
10 WEST 35TH STREET
CHICAGO, IL 60616
OICY ATTN TECHNICAL LIBRARY

INSTITUTE FOR DEFENSE ANALYSES
400 ARMY-NAVY DRIVE
ARLINGTON, VA 22202
OICY ATTN ERNEST BAUER
OICY ATTN HANS WOLFHARD
OICY ATTN JOEL BENGSTON
OICY ATTN PHILIP A SELWYN

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

TRT CORPORATION

P.O. BOX 81087

SAN DIEGO, CA 92138

OICY ATTN D A VROOM

OICY ATTN J A RUTHERFORD

OICY ATTN R H NEYNABER

JAYCOR

1401 CAMINO DEL MAR

DEL MAR, CA 92014

OICY ATTN H B LEVINE

JOHNS HOPKINS UNIVERSITY

APPLIED PHYSICS LABORATORY

JOHNS HOPKINS ROAD

LAUREL MD 20810

OICY ATTN DOCUMENT LIBRARIAN

JOHNS HOPKINS UNIVERSITY

34TH & CHARLES STREETS

BALTIMORE, MD 21218

OICY ATTN JOYCE J KAUFMAN

KAMAN SCIENCES CORPORATION

P. O. BOX 7463

COLORADO SPRINGS, CO 80933

OICY ATTN FRANK H SHELTON

OICY ATTN R J BITTNER

OICY ATTN P JESSOM

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

KENT STATE UNIVERSITY

DEPT OF CHEMISTRY

KENT OH 44242

OICY ATTN W C FERNELIUS

LOCKHEED MISSILES AND SPACE COMPANY
3251 HANOVER STREET

PALO ALTO, CA 94304

OICY ATTN BILLY M MCCORMAC DEPT 52-54

OICY ATTN MARTIN WALT DEPT 52-10

OICY ATTN RICHARD G JOHNSON DEPT 52-12

OICY ATTN TOM JAMES

OICY ATTN ROBERT C GUNTON DEPT 52-14

OICY ATTN ROBERT D SEARS DEPT 52-14

OICY ATTN J B REAGAN D/52-12

OICY ATTN JOHN KUMER

OICY ATTN JOHN L KULANDER DEPT 52-14

LOWELL UNIVERSITY OF

CENTER FOR ATMOSPHERIC RESEARCH

450 AIKEN STREET

LOWELL, MA 01854

OICY ATTN G T REST

MARYLAND UNIVERSITY OF

ADMINISTRATION BUILDING

COLLEGE PARK, MD 21030

OICY ATTN DEPT OF PHYSICS H GRIEM

OICY ATTN J VANDERSLICE CHEM DEPT

OICY ATTN ROBERT J MUHN DEPT OF CHEM

OICY ATTN T J ROSENBERG

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

MAXWELL LABORATORIES, INC.
9244 BALBOA AVENUE
SAN DIEGO, CA 92123
OICY ATTN VICTOR FARGO

MCDONNELL DOUGLAS CORPORATION
5301 BOLSA AVENUE
HUNTINGTON BEACH, CA 92647
OICY ATTN A D GOEDEKE
OICY ATTN WILLIAM OLSEN
OICY ATTN J MOILE

MIDWEST RESEARCH INSTITUTE
425 VOLKER BOULEVARD
KANSAS CITY, MO 64110
OICY ATTN PHYSICS SEC FRANK T GREENE
OICY ATTN PHYSICS SEC THOMAS H MILNE

MINNESOTA, UNIVERSITY OF
2030 UNIVERSITY AVENUE, S.F.
MINNEAPOLIS, MN 55414
(NO CLASSIFIED TO THIS ADDRESS)
OICY ATTN J R WINKLER

MINNESOTA, UNIVERSITY OF
MORRIS CAMPUS
MORRIS, MN 56267
OICY ATTN MERIE HIRSCH

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

MISSION RESEARCH CORPORATION
735 STATE STREET

SANTA BARBARA, CA 93101

01CY ATTN DAVE SOWLE

01CY ATTN RALPH KYLR

01CY ATTN W F CREVIER

01CY ATTN CONRAD L LONGMIRE

01CY ATTN NEAL J CARRON

01CY ATTN R HENDRICK

01CY ATTN R N WILSON

01CY ATTN R BOGUSCH

01CY ATTN M SCHEIRE

01CY ATTN D ARCHER

01CY ATTN D SAPPENFELD

01CY ATTN P FISCHER

MITRE CORPORATION, THE

P.O. BOX 208

REDFORD, MA 01730

01CY ATTN S A MORIN M/S

01CY ATTN J N FREEDMAN

NATIONAL ACADEMY OF SCIENCES

ATTN: NATIONAL MATERIALS ADVISORY BOARD

2101 CONSTITUTION AVENUE

WASHINGTON, DC 20418

02CY ATTN EDWARD R DYER

01CY ATTN J R STEVERS

01CY ATTN WILLIAM C BARTLEY

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

NEW YORK AT BUFFALO, STATE UNIV OF
BUFFALO, NY 14214

OICY ATTN GILBERT D BRINK

PACIFIC-SIERRA RESEARCH CORP.
1456 CLOVERFIELD BLVD.
SANTA MONICA, CA 90404
OICY ATTN F C FIELD JR

PENNSYLVANIA STATE UNIVERSITY
INDUSTRIAL SECURITY OFFICE
ROOM 5, OLD MAIN BUILDING
UNIVERSITY PARK, PA 16802
OICY ATTN L HALE
OICY ATTN J S NISBET

PHOTOMETRICS, INC.
442 MARRETT ROAD
LEXINGTON, MA 02173
OICY ATTN IRVING L KOFSKY

PHYSICAL DYNAMICS INC.
P.O. BOX 1069
BERKELEY, CA 94701
OICY ATTN A THOMPSON
OICY ATTN JOSEPH R WORKMAN

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

PHYSICAL SCIENCES, INC.
30 COMMERCE WAY
WOBURN, MA 01801
OICY ATTN KURT WRAY
OICY ATTN R L TAYLOR

PHYSICS INTERNATIONAL COMPANY
2700 MERCED STREET
SAN LEANDRO, CA 94577
OICY ATTN DDC CON FOR TECH LIR

PITTSBURGH, UNIVERSITY OF
OF THE COMWLTH SYS OF HIGHER EDUC
CATHEDRAL OF LEARNING
PITTSBURGH, PA 15213
OICY ATTN WADE L FITE
OICY ATTN MANFRED A RIONDI
OICY ATTN FREDERICK KAUFMAN
OICY ATTN EDWARD GERJUOY

PRINCETON UNIV, THE TRUSTEES OF
FORRESTAL CAMPUS LIBRARY
BOX 710
PRINCETON UNIVERSITY
PRINCETON, NJ 08540
(USE P.O. BOX 36 FOR UNCLASSIFIED ONLY)
OICY ATTN ARNOLD J KELLY

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

R & D ASSOCIATES

P.O. BOX 9695

MARINA DEL REY CA 90291

01CY ATTN R P TURCO

01CY ATTN RICHARD LATFER

01CY ATTN ROBERT E LELEVIER

01CY ATTN D DEE

01CY ATTN R G LINDGREN

01CY ATTN CHARLES H HUMPHREY

01CY ATTN BRYAN GARRARD

01CY ATTN H A DRY

01CY ATTN FORREST GILMORE

01CY ATTN ALBERT L LATFER

R & D ASSOCIATES

1815 N. FT. MYER DRIVE

11TH FLOOR

ARLINGTON, VA 22209

01CY ATTN HERRERY J MITCHELL

01CY ATTN J W ROSENGREN

RAND CORPORATION, THE

1700 MAIN STREET

SANTA MONICA, CA 90406

01CY ATTN CULLEN CRAIN

RICE UNIVERSITY

DEPARTMENT OF SPACE SCIENCE

HOUSTON, TX 77058

01CY ATTN RONALD F STERRINGS

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

RICE UNIVERSITY, WILLIAM MARSH

P.O. BOX 2692

HOUSTON, TX 77001

(ALL CORRES ATTN: INDUSTRIAL SEC SUPER)

OICY ATTN J CHAMBERLAIN

RIVERSIDE RESEARCH INSTITUTE

80 WEST END AVENUE

NEW YORK, NY 10023

OICY ATTN J R MINKOFF

OICY ATTN G GLASER

OICY ATTN DONALD H KOPPEL

SCIENCE APPLICATIONS, INC.

P.O. BOX 2351

LA JOLLA, CA 92038

OICY ATTN DANIEL A HAMLYN

OICY ATTN ROBERT W LOWEN

OICY ATTN BEN MYERS

SCIENCE APPLICATIONS, INC.

HUNTSVILLE DIVISION

2109 W. CLINTON AVENUE

SUITE 700

HUNTSVILLE, AL 35805

OICY ATTN NDEI R RYRN

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

SECHRIST, PROFESSOR CHALMERS F.
155 ELECTRICAL ENGINEERING BLDG.
UNIVERSITY OF ILLINOIS
URBANA, IL 61801
01CY ATTN C SECHRIST

SOUTHERN CALIFORNIA, UNIV OF
UNIVERSITY PARK
LOS ANGELES, CA 90007
01CY ATTN S W BENSON

STANFORD RESEARCH INSTITUTE
333 RAVENSWOOD AVENUE
MENLO PARK, CA 94025

01CY ATTN FELIX T SMITH (UNCL ONLY)
01CY ATTN D J HILDEBRAND
01CY ATTN L L CORR
01CY ATTN G BLACK
01CY ATTN J T MOSELEY
01CY ATTN RURT R GASTEN
01CY ATTN J LOMAX
01CY ATTN R D HAKE JR
02CY ATTN WALTER G CHESTNUT
01CY ATTN JAMES R PETERSON
01CY ATTN ARTHUR LEE WHITSON
01CY ATTN RAY L LEADABRAND
01CY ATTN ALLEN M PETERSON
01CY ATTN T G SLANGER
01CY ATTN RONALD WHITE

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

STANFORD RESEARCH INSTITUTE
1611 NORTH KENT STREET
ARLINGTON, VA 22209
01CY ATTN WARREN W BERNING

STANFORD RESEARCH INSTITUTE
306 WYNN DRIVE, N. W.
HUNTSVILLE, AL 35805
01CY ATTN MACPHERSON MORGAN

SYSTEMS, SCIENCE AND SOFTWARE, INC.
P.O. BOX 1620
LA JOLLA, CA 92038
01CY ATTN RALPH C SKLARFW

TECHNOLOGY INTERNATIONAL CORPORATION
75 WIGGINS AVENUE
BEDFORD, MA 01730
01CY ATTN W P ROQUITST

TRW SYSTEMS GROUP
ONE SPACE PARK
REDONDO BEACH, CA 90278
01CY ATTN R WATSON R1/1096
01CY ATTN J F FRICHTENTICHT R1-1196
01CY ATTN R K PLERUCH
01CY ATTN TECH INFO CENTER/S-1930

083

DATE 1/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

UNITED TECHNOLOGIES CORPORATION

755 MAIN STREET

HARTFORD, CT 06103

OICY ATTN ROBERT H BULLYS

OICY ATTN H MICHELS

UTAH STATE UNIVERSITY

LOGAN, UT 84321

OICY ATTN WILLIAM M MORE CHEMISTRY DEPT

OICY ATTN KAY BAKER

OICY ATTN C WYATT

OICY ATTN D BURT

OICY ATTN DORAN BAKER

VISIDYNE, INC.

19 THIRD AVENUE

NORTH WEST INDUSTRIAL PARK

BURLINGTON, MA 01803

OICY ATTN T C DEGGES

OICY ATTN J W CARPENTER

OICY ATTN OSCAR MANLEY

OICY ATTN R WALTZ

WAYNE STATE UNIVERSITY

1064 MACKENZIE HALL

DETROIT, MI 48202

OICY ATTN R H KIMMLER

OICY ATTN PIETER K ROL CHEM ENGRG & MAT SCI

083

DATE 23/02/77

DEPARTMENT OF DEFENSE CONTRACTORS

WAYNE STATE UNIVERSITY
DEPT. OF PHYSICS
DETROIT, MI 48202

01CY ATTN WALTER F KAUPPILA

YALE UNIVERSITY
NEW HAVEN, CT 06520

01CY ATTN ENGRG DEPT GEORGE J SCHULTZ

ED
77